

Problem Set 4

Due: Week 11

Multiple choice questions (2 points each)

1. Which of the following defines a Geometric distribution?
 - (a) It models the number of successes in a fixed number of independent trials.
 - (b) It models the number of trials needed to get the first success in a sequence of independent Bernoulli trials.
 - (c) It models the number of events occurring in a fixed interval of time.
 - (d) It models the number of successes in a Poisson process.
2. If X is a Poisson random variable with parameter $\lambda = 3$, what is $E(X)$?
 - (a) 1
 - (b) 3
 - (c) 6
 - (d) 9
3. In a factory, 10% of the products produced are defective. Let X be the random variable representing the number of defective products in a batch of 20 products. Which of the following distributions best models this scenario?
 - (a) $X \sim \text{Bernoulli}(p = 0.1)$
 - (b) $X \sim \text{Binomial}(n = 20, p = 0.1)$
 - (c) $X \sim \text{Geometric}(p = 0.1)$
 - (d) None of the above
4. A call center receives an average of 5 calls per hour. Let X be the random variable representing the number of calls received in the next hour. Which of the following distributions best models this scenario?
 - (a) $X \sim \text{Bernoulli}(p = 5)$
 - (b) $X \sim \text{Binomial}(n = 7, p = 5/60)$
 - (c) $X \sim \text{Poisson}(\lambda = 5)$
 - (d) None of the above

5. Suppose that a certain type of magnetic tape contains on average three defects per 1000 feet. What is the probability that a roll of tape 1200 feet long contains no defects?
- $1 - \frac{3}{1000}$
 - $(1 - \frac{3}{1000})^{\frac{1200}{1000}}$
 - $e^{-3.6}$
 - None of the above
6. Which of the following statements about joint probability distributions is true?
- They describe the probability of two or more independent happened at the same time
 - They are composed from the probability distribution of each random variable
 - They describe the probability of two or more random variables occurring simultaneously
 - None of the above
7. Given the joint distribution $f(x, y)$ and marginal distributions $f_X(x)$ and $f_Y(y)$. Which of the following is true?
- $f_X(x)$ is a probability function only about random variable X
 - $f_X(x) = P(X = x|Y = y)$
 - $f(x, y) = f_X(x)f_Y(y)$
 - $f(x, y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_X(x)f_Y(y)dx dy$
8. If $f(x, y)$ represents the joint probability distribution of two discrete random variables X and Y , which of the following expressions correctly computes the marginal probability function $f_X(x)$?
- $f_X(x) = f(x, y_0)$ for some specific y_0
 - $f_X(x) = \sum_y f(x, y)$
 - $f_X(x) = f(x|y)f(y)$
 - $f_X(x) = \int_{-\infty}^{\infty} f(x, y)dy$
9. Suppose that X and Y have a discrete joint distribution for which the joint PMF is defined as follows:

$$p(x, y) = \begin{cases} c|x + y| & \text{for } x = -2, -1, 0, 1, 2 \text{ and} \\ & y = -2, -1, 0, 1, 2 \\ 0 & \text{otherwise} \end{cases}$$

- (3 points) Determine the value of c .
- (3 points) Find the marginal distribution of X .
- (4 points) Find $P(|X - Y| \leq 1)$